

PFA AND IDEALS ON ω_2 WHOSE ASSOCIATED FORCINGS ARE PROPER

SEAN COX

ABSTRACT. Given an ideal I , let $\mathbb{P}_I$ denote the forcing with I -positive sets. We consider models of forcing axioms $MA(\Gamma)$ which also have a normal ideal I with completeness ω_2 such that $\mathbb{P}_I \in \Gamma$. Using a bit more than a superhuge cardinal, we produce a model of PFA which has many ideals on ω_2 whose associated forcings are proper; a similar phenomenon is also observed in the standard model of $MA^{+\omega_1}(\sigma\text{-closed})$ obtained from a supercompact cardinal. Our model of PFA also exhibits weaker versions of ideal properties which were shown in [9] to be inconsistent with PFA.

Along the way, we also show (1) the Diagonal Reflection Principle for internally club sets ($DRP(IC_{\omega_1})$) introduced in [4] is equivalent to a natural weakening of “There is an ideal I such that $\mathbb{P}_I$ is proper”; and (2) For many natural classes Γ of posets, $MA^{+\omega_1}(\Gamma)$ is equivalent to an apparently stronger version which we call $MA^{+Diag}(\Gamma)$.

1. INTRODUCTION

In [4], we introduced the *Diagonal Reflection Principle (DRP)*—which is a highly simultaneous form of stationary set reflection—and proved that DRP follows from strong forcing axioms. DRP asserts that a certain naturally-defined set of ω_1 -sized structures—namely, those structures on which the nonstationary ideal condenses correctly—is *stationary* (see section 3 below; such condensation notions originally appeared in Foreman [8]). Independently, Viale [13] proved similar results.

A motivation for the paper is the following observation: if F is a filter which concentrates on sets witnessing DRP , then by examining generic ultrapowers which use F (such ultrapowers have critical point ω_2), we

2000 *Mathematics Subject Classification.* 03.

Key words and phrases. Ideals, Forcing Axioms, PFA.

I thank Matt Foreman and Menachem Magidor for helpful conversations on related topics.

see that $\mathbb{P}_{\check{F}}$ exhibits properties which resemble *properness* (This observation follows from Theorem 8 below). So a natural way to strengthen DRP is to require that $\mathbb{P}_{\check{F}}$ is actually proper.

Most of the paper is devoted to models of strong forcing axioms $MA(\Gamma)$ which have an ideal I such that $\mathbb{P}_I \in \Gamma$; in particular, we obtain such ideals whose dual filter concentrates on sets witnessing DRP. In Section 4 we observe that, in the standard model of $MA^{+\omega_1}(\sigma\text{-closed})$ obtained from collapsing everything below a supercompact cardinal to ω_1 , there are ideals whose duals concentrate on sets witnessing DRP and whose associated forcings have σ -closed dense subsets. It is natural to ask if a similar situation can arise under PFA; we answer this question affirmatively. In particular, we prove the following theorem, which is stated more precisely as Theorem 12 in Section 5:

Theorem. *Relative to a super-2-huge cardinal, it is consistent that PFA holds and there are ideals I such that $\mathbb{P}_I$ is proper. Moreover, there are such ideals whose dual concentrates on sets witnessing DRP.*

This model has several other interesting features. In [9], Foreman and Magidor showed that if *either*:

- (1)
 - $(\omega_3, \omega_2) \twoheadrightarrow (\omega_2, \omega_1)$; or
 - There is a presaturated ideal on ω_2

then PFA fails. On the other hand, the model of PFA we produce in Section 5 shows that weaker versions of (1)—namely that $(\theta, \omega_2) \twoheadrightarrow (\omega_2, \omega_1)$ holds for an inaccessible θ and that there is an ideal on ω_2 with closure properties resembling, but weaker than, presaturation—are (simultaneously) consistent with PFA.

Finally, it is well-known that PFA implies every proper poset $\mathbb{Q}$ completely embeds below some condition in $[H_\theta]^{\omega_1}/NS$ (for all sufficiently large θ); this is due to the existence of stationarily many $M \in \wp_{\omega_2}(H_\theta)$ such that $\omega_1 \subset M$ and there exists an $(M, \mathbb{Q})$ -generic; let $S_{\mathbb{Q}}^{H_\theta}$ denote this stationary set.¹ So in particular, if PFA holds and I is an ideal whose associated forcing $\mathbb{P}_I$ is proper, then $\mathbb{P}_I$ completely embeds into another ideal forcing $\mathbb{P}_{I'}$ where I' concentrates on $S_{\mathbb{P}_I}^{H_{\theta'}}$ (namely I' is the nonstationary ideal restricted to $S_{\mathbb{P}_I}^{H_{\theta'}}$). In general the nature of this complete embedding $\mathbb{P}_I \rightarrow \mathbb{P}_{I'}$ is mysterious; however we produce a model of PFA with such complete embeddings of a simple form (the proof appears in Section 6):

Theorem. *Relative to a super-3-huge cardinal, it is consistent that PFA holds and there are ideals I, I' such that $\mathbb{P}_I, \mathbb{P}_{I'}$ are proper, I'*

¹This is a general fact about forcing axioms, not just PFA; see Section 2.

projects to I in the Rudin-Keisler sense, and this projection is also a projection in the sense of forcing.

The paper is organized as follows: Section 2 gives background material and notation, including a discussion “plus” versions of forcing axioms and Foreman’s Duality Theorem (which is heavily used throughout the paper); Section 3 discusses DRP and characterizations in terms of ideals on ω_2 ; Section 4 is about models of $MA^{+\omega_1}(\sigma\text{-closed})$ with σ -closed ideal forcings; Section 5 is about models of PFA with proper ideal forcings; and Section 6 discusses “Ideal projections as forcing projections” in models of PFA.

2. NOTATION AND BACKGROUND

2.1. Ideals. $\wp_\kappa(H_\theta)$ denotes the set of $M \prec H_\theta$ such that $|M| < \kappa$ and $M \cap \kappa \in \kappa$. If Z is a set, $NS \restriction Z$ denotes the ideal $\{A \subset Z \mid A \text{ is nonstationary in } \bigcup Z\}$.² Throughout this paper, *ideal* always means a normal ideal. If I is an ideal then $\check{I}$ denotes the filter which is dual to I ; similarly if F is a filter then $\check{F}$ denotes its dual ideal. I^+ denotes the I -positive sets (i.e. if $I \subset \wp(Z)$ then I^+ is the collection of $S \in \wp(Z)$ such that $S \notin I$); if F is a filter then F^+ means $\check{F}^+$. If Δ is a class and F a filter, we say that F *concentrates on* Δ iff there is some $A \in F$ such that $A \subseteq \Delta$ (and an ideal I concentrates on Δ iff $\check{I}$ concentrates on Δ). The *forcing associated with* I is $(I^+, \subset)$, which we will denote $\mathbb{P}_I$; this is equivalent to forcing with $\wp(Z)/I - \{0\}$, where by definition $S =_I T$ iff the symmetric difference of S and T is in I .

$(\theta, \mu) \twoheadrightarrow (\theta', \mu')$ denotes the statement: For every first order structure $\mathcal{A}$ on θ , there is a $Z \prec \mathcal{A}$ such that $|Z| = \theta'$ and $|Z \cap \mu| = \mu'$ (the classic *Chang’s Conjecture* is the special case $(\omega_2, \omega_1) \twoheadrightarrow (\omega_1, \omega)$).

We refer to a notion related to saturation of an ideal; this notion is analyzed in more detail in [3], though no results from that paper are used here.

Definition 1. Let H be a set (typically $\theta \subseteq H \subseteq H_\theta$ for some θ), $Z \subset \wp(H)$, and $I \subset \wp(Z)$ be an ideal. Suppose I' is an ideal over some $Z' \subset H_{\theta'}$ for $\theta' \gg |H|$ such that I' projects to I ; i.e. $I = \{\{Z' \cap H \mid Z' \in A'\} \mid A' \in I'\}$; let $\pi : (I')^+ \rightarrow I^+$ be the map $S \mapsto \{M \cap H \mid M \in S\}$.

- If π is a projection in the sense of forcing (i.e. pointwise preimages of maximal antichains are maximal), then we say I, I' witnesses “Ideal projections as forcing projections” and write $FP(I', I)$.

²For example, if $Z = [H_\theta]^\omega$ then $A \in NS \restriction Z$ iff $A \subset [H_\theta]^\omega$ and there is some $F : [H_\theta]^{<\omega} \rightarrow H_\theta$ such that no element of A is closed under F .

- If there is some I' such that $FP(I', I)$ holds, then we say $FP(I)$ holds.
- If $FP(I', I)$ holds where I' is the conditional club filter relative to I at H_θ ,³ we say $FP^{ConditionalClub}(I)$ holds.

In [3] it is shown that $FP(I)$ implies precipitousness of I , and that $FP^{ConditionalClub}(I)$ is equivalent to saturation of I (the latter is essentially due to Lemma 3.46 of [7]). In the present paper we produce a model of $FP(I)$ where I is not saturated (or even presaturated).

2.2. Forcing Axioms. By $MA(\Gamma)$ we always mean $MA_{\omega_1}(\Gamma)$; i.e. for every $\mathbb{P} \in \Gamma$ and every ω_1 -sized collection $\mathcal{D}$ of dense subsets of $\mathbb{P}$ there is a filter on $\mathbb{P}$ which meets every set in $\mathcal{D}$. For an ordinal α , $MA^{+\alpha}(\Gamma)$ means that for every $\mathbb{P} \in \Gamma$, every ω_1 -sized collection $\mathcal{D}$ of dense subsets of $\mathbb{P}$, and every sequence $\langle \dot{S}_i | i < \alpha \rangle$ of $\mathbb{P}$ -names such that $\Vdash_{\mathbb{P}} \text{“}\dot{S}_i \text{ is stationary subset of } \omega_1\text{”}$ for every $i < \alpha$, there is a filter $F \subset \mathbb{P}$ which meets every D_i and for every $i < \omega_1$: $(\dot{S}_i)_F := \{\beta : \text{There is some } q \in F \text{ such that } q \Vdash \check{\beta} \in \dot{S}_i\}$ is stationary.

The proof of Theorem 2.53 in [14] shows that $MA(\{\mathbb{P}\})$ is equivalent to stationarity of the following set:

$$(2) \quad S_{\mathbb{P}}^{H_\theta} := \text{the set of } M \prec (H_\theta, \in, \{\mathbb{P}\}) \text{ such that } \omega_1 \subset M \text{ and there exists an } (M, \mathbb{P})\text{-generic object}$$

for all (some) sufficiently large θ . The same argument shows that $MA^{+\alpha}(\{\mathbb{P}\})$ is equivalent to the statement: for every sequence $\langle \dot{S}_i | i < \alpha \rangle$ of $\mathbb{P}$ -names for stationary subsets of ω_1 , the set:

$$(3) \quad S_{\dot{S}, \mathbb{P}}^{H_\theta} := \text{the set of } M \prec (H_\theta, \in, \{\mathbb{P}, \dot{S}\}) \text{ such that } \omega_1 \subset M \text{ and there exists a } g \text{ which is } (M, \mathbb{P})\text{-generic and } \dot{S}_g \text{ is stationary for every } i < \alpha.$$

is stationary for all (some) sufficiently large θ . Using the latter equivalence as motivation, let us define the following apparent strengthening of $MA^{+\omega_1}(\{\mathbb{P}\})$:

- Definition 2.** • For a poset $\mathbb{P}$ and a regular cardinal θ such that $\wp(\mathbb{P}) \in H_\theta$: $S_{\mathbb{P}}^{+Diag, H_\theta}$ will denote the set of $M \prec (H_\theta, \in, \{\mathbb{P}\})$ such that $\omega_1 \subset M$ and there exists a g which is $(M, \mathbb{P})$ -generic and such that $\dot{S}_g$ is stationary whenever $\dot{S} \in M$ is a $\mathbb{P}$ -name for a stationary subset of ω_1 .
- $MA^{+Diag}(\Gamma)$ denotes the statement that for all $\mathbb{P} \in \Gamma$ and for sufficiently large regular θ , $S_{\mathbb{P}}^{+Diag, H_\theta}$ is stationary.

³This is defined in [7]

There are obvious generalizations of Definition 2 which require stationary evaluation of names of stationary subsets of $[\lambda]^\omega$ by g .⁴ We note the following:

Lemma 3. *Suppose Γ is a class of posets such that whenever $\mathbb{P} \in \Gamma$ then $\mathbb{P} * \text{Col}(\omega_1, \wp(\omega_1)^{V[\dot{G}]}) \in \Gamma$ (where $\dot{G}$ is the canonical $\mathbb{P}$ -name for the $\mathbb{P}$ -generic object). Then $MA^{+\omega_1}(\Gamma)$ is equivalent to $MA^{+Diag}(\Gamma)$.*

Proof. To see that $MA^{+Diag}(\Gamma)$ implies $MA^{+\omega_1}(\Gamma)$: Fix any $\mathbb{P} \in \Gamma$ and a regular θ such that $\wp(\mathbb{P}) \in H_\theta$. If $\langle \dot{S}_i \mid i < \omega_1 \rangle$ is a sequence of $\mathbb{P}$ -names for stationary subsets of ω_1 , then almost every $M \in S_{\mathbb{P}}^{+Diag, H_\theta}$ has $\dot{S}$ as an element; since $\omega_1 \subset M$ then $\dot{S}_i \in M$ for each i . Then if g is any $(M, \mathbb{P})$ -generic which witnesses that $M \in S_{\mathbb{P}}^{+Diag, H_\theta}$, then $(\dot{S}_i)_g$ is stationary for each $i < \omega_1$.

Now assume $MA^{+\omega_1}(\Gamma)$ and let $\mathbb{P} \in \Gamma$; we want to show that $MA^{+Diag}(\mathbb{P})$ holds. Let $\dot{Q}$ be the $\mathbb{P}$ -name for $\text{Col}(\omega_1, \wp(\omega_1)^{V[\dot{G}]})$, where $\dot{G}$ is the canonical $\mathbb{P}$ -name for the $\mathbb{P}$ -generic; so by assumption, $\mathbb{P} * \dot{Q} \in \Gamma$. Let $\langle \dot{T}_\alpha \mid \alpha < \omega_1 \rangle$ be a $\mathbb{P} * \dot{Q}$ -name which is forced to enumerate $\{S \in V[G] \mid V[G] \models S \subseteq \omega_1 \text{ is stationary}\}$. Now for each $\alpha < \omega_1$, $\mathbb{P} * \dot{Q}$ forces that $\dot{T}_\alpha$ is stationary (since $\dot{Q}_G$ is countably closed and thus stationary set preserving in $V[G]$). Then by $MA^{+\omega_1}(\mathbb{P} * \dot{Q})$, $S_{\dot{T}, \mathbb{P} * \dot{Q}}^{H_\theta}$ is stationary. Let M be any element of $S_{\dot{T}, \mathbb{P} * \dot{Q}}^{H_\theta}$, let $\dot{S} \in M$ be a $\mathbb{P}$ -name for a stationary subset of ω_1 , and let $g * h$ be an $(M, \mathbb{P} * \dot{Q})$ -generic witnessing that $M \in S_{\dot{T}, \mathbb{P} * \dot{Q}}^{H_\theta}$. Then $\dot{S}_g = (\dot{T}_\alpha)_{g * h}$ for some $\alpha < \omega_1$, and the latter is stationary since $M \in S_{\dot{T}, \mathbb{P} * \dot{Q}}^{H_\theta}$. $\square$

Corollary 4. *$PFA^{+\omega_1}$ is equivalent to PFA^{+Diag} , $MM^{+\omega_1}$ is equivalent to MM^{+Diag} , and $MA^{+\omega_1}(\sigma\text{-closed})$ is equivalent to $MA^{+Diag}(\sigma\text{-closed})$.*

Proof. Let Γ be one of those 3 classes of posets (i.e. proper, stationary set preserving for subsets of ω_1 , or σ -closed). Then for every $\mathbb{P} \in \Gamma$, $\mathbb{P} * \text{Col}(\omega_1, \wp(\omega_1)^{V[\dot{G}]}) \in \Gamma$. Apply Lemma 3. $\square$

2.3. Diagonal Reflection. We recall the definition of DRP from [4]:

Definition 5. *Let Z be a class of ω_1 -sized sets (for example, Z could be the class of all ω -closed sets).*

The Diagonal Reflection Principle at θ relative to Z , abbreviated $DRP(\theta, Z)$, is the statement:

⁴if $\mathbb{P}$ satisfies certain requirements this often follows from $MA^{+Diag}(\mathbb{P})$ for sufficiently small λ ; for example, if $\mathbb{P}$ is σ -closed and $\Vdash_{\mathbb{P}} |H_\theta^V| = \omega_1$ then $MA^{+\omega_1}(\mathbb{P})$ implies this generalized version for all small λ .

There are stationarily many $M \in \wp_{\omega_2}(H_{(\theta^\omega)^+})$ such that:

- (1) $M \cap H_\theta \in Z$
- (2) Whenever $R \in M$ is a stationary subset of $[\theta]^\omega$, then $R \cap [M \cap \theta]^\omega$ is stationary in $[M \cap \theta]^\omega$.

We say *DRP holds on Z* iff $DRP(\theta, Z)$ holds for all regular $\theta \geq \omega_2$.

Definition 6. $wDRP(\theta, Z)$ is defined exactly like $DRP(\theta, Z)$ except that we replace clause 2 of the definition with the following:

- Whenever $R \in M$ is a **projective** stationary subset of $[\theta]^\omega$, then $R \cap [M \cap \theta]^\omega$ is stationary in $[M \cap \theta]^\omega$.

Independently, Viale [13] considered notions very similar to *DRP* and proved similar theorems to those in [4]. Adopting his terminology, for an ordinal λ of cofinality at least ω_2 and an $M \prec (H_\theta, \in, \{\lambda\} \dots)$, let us say M is $\lambda \cap \text{cof}(\omega)$ -*faithful* iff $R \cap \sup(M \cap \lambda)$ is stationary in $\sup(M \cap \lambda)$ for every $R \in M$ which is a stationary subset of $\lambda \cap \text{cof}(\omega)$. Similarly we will say M is $[\lambda]^\omega$ -*faithful* if the analogous statement holds for every $R \in M$ which is a stationary subset of $[\lambda]^\omega$.

For this paper, the most relevant classes Z in Definition 5 are the classes of *internally approachable* (IA_{ω_1}) and *internally club* (IC_{ω_1}) structures of size ω_1 . IA_{ω_1} is the class of all M such that there is some $\in$ -increasing and $\subset$ -continuous chain $\langle N_\xi \mid \xi < \omega_1 \rangle$ of countable sets such that $M = \bigcup_{\xi < \omega_1} N_\xi$ and every proper initial segment of $\vec{N}$ is an element of M . IC_{ω_1} is the (possibly wider) class defined similarly, except that we only require that each N_ξ is an element of M (equivalently, $M \in IC_{\omega_1}$ iff $M \cap [M]^\omega$ contains a club).

2.4. Forcing quotients and Foreman's Duality Theorem. Following [7], if $\mathbb{P}$ and $\mathbb{Q}$ are posets, then $i : \mathbb{P} \rightarrow \mathbb{Q}$ is a *regular embedding of $\mathbb{P}$ to $\mathbb{Q}$* iff i preserves the order, preserves incompatibility, and pointwise maps maximal antichains in $\mathbb{P}$ to maximal antichains in $\mathbb{Q}$. If $i : \mathbb{P} \rightarrow \mathbb{Q}$ is regular and $G \subset \mathbb{P}$ is generic, then $\mathbb{Q}/i''G$ denotes the collection of $q \in \mathbb{Q}$ such that q is compatible with every member of $i''G$; the ordering on $\mathbb{Q}/i''G$ is just the ordering inherited from $\mathbb{Q}$. Regularity of i ensures that $\mathbb{Q}$ is forcing equivalent to $\mathbb{P} * \mathbb{Q}/i''\dot{G}$ (where $\dot{G}$ is the canonical $\mathbb{P}$ -name for its generic object).

We will often use the following construction of precipitous filters, and caution that it varies slightly from many of the constructions in [7] because the construction below yields a filter which does *not* extend

the ground model's ultrafilter.⁵ Suppose $j : V \rightarrow_U N$ is an ultrapower embedding via some normal ultrafilter $U \in V$ which concentrates on $\{M \mid M \prec H_\lambda\}$ and has critical point κ (where $\kappa \leq \lambda$). Let $\mathbb{P} \in H_\lambda$ be a poset; note $j \restriction \mathbb{P} : \mathbb{P} \rightarrow j(\mathbb{P})$ preserves order and incompatibility. Assume also that:

(4) $j \restriction \mathbb{P}$ is a regular embedding from $\mathbb{P}$ to $j(\mathbb{P})$

This happens, for instance, whenever $\mathbb{P}$ has the κ -cc, which will always be the case in this paper.⁶ So $j(\mathbb{P})$ is forcing equivalent to $\mathbb{P} * j(\mathbb{P})/j''\dot{G}$. Then whenever G is $(V, \mathbb{P})$ -generic and H is $(V[G], j(\mathbb{P})/j''\dot{G})$ -generic, in $V[G][H]$ the map j can be lifted to a $\hat{j}_{G*H} : V[G] \rightarrow N[G][H]$ via $\tau_G \mapsto j(\tau)_{H'}$, where H' is the $(V, \mathbb{P} * j(\mathbb{P})/j''\dot{G})$ -generic obtained by transferring $G * H$ via the forcing equivalence of $\mathbb{P} * j(\mathbb{P})/j''\dot{G}$ with $j(\mathbb{P})$. This map will be well-defined and elementary (see Section 9 of [5] for details). Also, $\hat{j}_{G*H}(G)$ is equal to the generic for $(V, j(\mathbb{P}))$ obtained by transferring $G * H$ to a generic for $j(\mathbb{P})$ modulo the equivalence of $j(\mathbb{P})$ with $\mathbb{P} * j(\mathbb{P})/j''\dot{G}$. This last fact can be used to show that $N \cap \hat{j}_{G*H}''H_\lambda[G] = j''H_\lambda$. Now $\hat{j}_{G*H}''H_\lambda[G]$ is an element of $N[G][H]$ and

(5) $U_{G*H} := \{A \in V[G] \mid A \subset \wp(H_\lambda[G]) \text{ and } \hat{j}_{G*H}''H_\lambda[G] \in \hat{j}_{G*H}(A)\}$

is a $V[G]$ -normal ultrafilter.⁷ We caution again that we are using $\hat{j}_{G*H}''H_\lambda[G]$, not $j''H_\lambda$, to define the ultrafilter U_{G*H} and as a result U_{G*H} will typically *not* extend U (e.g. if $\mathbb{P}$ turns κ into a successor cardinal, then U_{G*H} concentrates on models which believe κ is a successor cardinal, while U concentrates on models which believe κ is inaccessible). Still, using the definition of U_{G*H} , the fact that $N \cap \hat{j}_{G*H}''H_\lambda[G] = j''H_\lambda$, and the assumption that j is an ultrapower embedding which maps $\mathbb{P}$ regularly into $j(\mathbb{P})$, then U_{G*H} will always concentrate on the set A of those $M' \prec H_\lambda[G]$ such that:

- $M' \cap V \in V$; let M denote $M' \cap V$;
- M is an element of the underlying set $\bigcup \bigcup U$ measured by U (e.g. if U is an ultrafilter on $\wp(S)$ then $M' \cap V \in S$).

⁵The reason for this is that we want the filter to concentrate on elementary substructures of the generic extension, and thus cannot hope that a measure one set of such structures has any intersection at all with the ground model.

⁶If $A \subset \mathbb{P}$ is a maximal antichain, then $M \models "j(A) \text{ is a maximal antichain in } j(\mathbb{P})"$ and this clearly then holds in V as well. Since $|A| < \kappa = \text{cr}(j)$, $j(A) = j[A]$; so $j[A]$ is maximal in $j(\mathbb{P})$.

⁷Whenever $j : M \rightarrow N$ is an embedding (not necessarily definable in M), $d \in M$, and $j''d \in N$, then the ultrafilter $\{A \in M \mid j''d \in j(A)\}$ will always be normal with respect to M .

- $V \models "M \cap \mathbb{P} \text{ is a regular subposet of } \mathbb{P}"$ (Recall we are assuming (4)).

Using these facts, it can be shown that $N[G][H] = \{\hat{j}_{G*H}(f)(\hat{j}_{G*H}'' H_\lambda[G]) \mid f \in V[G] \cap {}^A V[G]\}$ (recall from above that $A \in V[G]$ will always have measure one in U_{G*H}) and it follows that $\hat{j}_{G*H}$ is the ultrapower map corresponding to $ult(V[G], U_{G*H})$.

In $V[G]$ define the following functions on A (here we again use the notation $M := M' \cap V$ whenever $M' \cap V \in V$):

- (1) $Q(M') := \mathbb{P}/(G \cap M)$
- (2) $h(M') :=$ the generic for $\mathbb{P}/(G \cap M)$ obtained by G and the forcing equivalence between $\mathbb{P}$ and $(\mathbb{P} \cap M) * \mathbb{P}/(\dot{G} \cap \dot{M})$
- (3) Given a $q \in j(\mathbb{P})/j''G$, note that $q \in V$ and that there is some $f_q \in V$ such that $q = [f_q]_U$ (here U was the ultrafilter in the ground model). Then define in $V[G]$ the function f'_q on A by $M' \mapsto f_q(M)$.

Then one can check:

For any H which is $(V[G], j(\mathbb{P})/j''G)$ -generic:

- (6)
 - (1) $[Q]_{U_{G*H}} = j(\mathbb{P})/j''G$;
 - (2) $[h]_{U_{G*H}} = H$;
 - (3) For every $q \in j(\mathbb{P})/j''G$, $[f'_q]_{U_{G*H}} = q$.

Definition 7. In $V[G]$, $F(j)$ denotes the collection of all $A \subset {}^\omega(H_\lambda)$ such that $\Vdash_{j(\mathbb{P})/j''G} \dot{A} \in \dot{U}_{G*\dot{H}}$.

$F(j)$ is a filter in $V[G]$ and inherits the completeness and normality properties from U (see Section 3.2 of [7]). By (6) and a special case of Foreman's Duality Theorem—namely Proposition 7.13 of [7]—we have:

- (7) In $V[G]$, the map $A \mapsto \|\dot{A} \in \dot{U}_{G*\dot{H}}\|_{ro(j(\mathbb{P})/j''G)}$ is an isomorphism between $F(j)^+$ and a dense subset of $ro(j(\mathbb{P})/j''G)$.

3. EQUIVALENT FORMULATIONS OF DRP, AND EFFECT OF DRP ON GENERIC EMBEDDINGS

In this section we show that the existence of ideals whose forcings are proper implies DRP (at least if the ideal concentrates on IC_{ω_1}), and in turn DRP always yields ideals whose associated forcings resemble proper forcings in a weak sense. Namely, DRP implies that for sufficiently small λ , stationary subsets of $[\lambda]^\omega$ are not destroyed in the generic ultrapower of V (though they may be destroyed in the generic extension!). In [4] we showed the following (see Section 2.3 for the definition of IC_{ω_1} and IA_{ω_1}):

- $MA^{+\omega_1}(\sigma\text{-closed})$ implies $DRP(\theta, IA_{\omega_1})$ for all regular $\theta \geq \omega_2$ (and MM implies $wDRP(\theta, IA_{\omega_1})$ for every regular $\theta \geq \omega_2$; see Definition 6).
- $DRP(\theta, IC_{\omega_1})$ implies that there is a stationary set of $M \in IC_{\omega_1} \cap \wp_{\omega_2}(H_{(\theta^\omega)^+})$ on which $NS \restriction [H_\theta]^\omega$ *condenses correctly*; i.e. such that if $\sigma : H_M \rightarrow M$ is the inverse of the transitive collapse of M and $\bar{H} := \sigma^{-1}(H_\theta)$, then $(NS \restriction [\bar{H}]^\omega)^{H_M}$ coheres with $(NS \restriction [\bar{H}]^\omega)^V$.

There are several equivalent formulations of $DRP(\theta, IC_{\omega_1})$; note formulations C and D below look like weak versions of saying $\mathbb{P}_I$ is proper.

Theorem 8. *For regular $\theta \geq \omega_2$, let Z^θ denote the collection of $M \in \wp_{\omega_2}(H_{(\theta^\omega)^+})$ such that $M \cap H_\theta \in IC_{\omega_1}$. The following are equivalent:*

- (A) $DRP(\theta, Z^\theta)$
- (B) *There are stationarily many $M \in Z^\theta$ such that $NS \restriction [H_\theta]^\omega$ condenses correctly via M*
- (C) *There is a normal ideal I whose dual contains the club filter over Z^θ such that $\Vdash_{\mathbb{P}_I}$ “every stationary $R \subset [H_\theta]^\omega$ from the ground model remains stationary in $ult(V, \dot{G})$ ”;*
- (D) *There is a stationary $S \subset Z^\theta$ such that $S \Vdash_{NS}$ “every stationary $R \subset [H_\theta]^\omega$ from the ground model remains stationary in $ult(V, \dot{G})$.”*

Proof. Note that if $M \cap H_\theta \in IC_{\omega_1}$, then $M \cap [M \cap H_\theta]^\omega$ contains a club.⁸ This implies that, letting $\sigma_M : H_M \rightarrow M$ be the inverse of the Mostowski Collapse map and $\bar{H} := \sigma_M^{-1}(H_\theta)$, if $S \in M$ is a stationary subset of $[H_\theta]^\omega$, then $S \cap [M \cap H_\theta]^\omega$ is stationary if and only if V believes that $\sigma_M^{-1}(S)$ is stationary in $[\bar{H}]^\omega$ (see the proof of Theorem 3.6 of [4] for more details). This gives the equivalence of (A) with (B).

For the remainder of the proof, we use the standard fact⁹ that if U is a normal V -ultrafilter on some $Z \subset \wp_{\omega_2}(H_\Gamma)$ (e.g. if U is generic for the forcing with a normal ideal) and $j : V \rightarrow_U ult(V, U)$ is the ultrapower, then:

$$(8) \quad \begin{array}{l} j \restriction H_\Gamma \in ult(V, U) \text{ and is represented by } [M \mapsto \sigma_M]_U, \\ \text{where } \sigma_M \text{ is the inverse of the Mostowski Collapse of} \\ M. \end{array}$$

⁸Thus the name; in fact this characterizes IC_{ω_1} .

⁹This is essentially Claim 2.26 of [7], though that claim is specifically about generic ultrapowers by normal ideals. Roughly, normality of U with respect to functions from V ensures that the $\in_U$ -extension of $[id]_U$ is equal to $j_U'' H_\Gamma$. This, in turn, implies that the transitive collapse of $[id]_U$ as computed in $ult(V, U)$ is in the wellfounded part of $ult(V, U)$. Assuming the wellfounded part of $ult(V, U)$ has been transitivity, this transitive collapse is equal to H_Γ and the inverse of the collapsing map is $j \restriction H_\Gamma$.

To see that (A) implies (D): Let S be the stationary set which witnesses $DRP(\theta, Z^\theta)$ and G be generic for $(V, \wp(Z^\theta)/(NS \upharpoonright Z^\theta))$ with $S \in G$, and $j : V \rightarrow_G ult(V, G)$ be the generic ultrapower. Let $R \in V$ be a stationary subset of $[H_\theta]^\omega$; note by the definition of Z^θ , there are G -many models with $[H_\theta]^\omega$ and R as elements. By (8), together with Los' Theorem and the fact that (A) implies (B), $ult(V, G)$ believes that $NS \upharpoonright [j(H_\theta)]^\omega$ condenses correctly via j^* , where $j^* := j \upharpoonright H_{(\theta^\omega)^+}$. In particular, $ult(V, G)$ believes that $(j^*)^{-1}(j(R)) = R$ is stationary.

(D) clearly implies (C).

Finally, we prove that (C) implies (A). Let G be generic for $(V, \mathbb{P}_I)$ and $j : V \rightarrow_G ult(V, G)$ be the generic embedding. Then for every stationary $R \subset [H_\theta]^\omega$: since R remains stationary in $ult(V, G)$, and since $ult(V, G) \models$ “ $j''H_{(\theta^\omega)^+} \cap j(H_\theta)$ is internally club,” then $ult(V, G) \models$ “ $j(R)$ reflects to $j''H_\theta$.” So by Los' Theorem there are G -many structures M such that every $R \in M$ which is a stationary subset of $[H_\theta]^\omega$ reflects to $M \cap H_\theta$. Since the dual of I extends the club filter, this collection of M is stationary and witnesses $DRP(\theta, Z^\theta)$. $\square$

Corollary 9. *Suppose I is a normal ideal which concentrates on $IC_{\omega_1} \cap \wp_{\omega_2}(H_{(\theta^\omega)^+})$ such that $\mathbb{P}_I$ is proper. Then $DRP(\theta, IC_{\omega_1})$ holds.*

Proof. This is an immediate corollary of part C of Theorem 8, the definition of properness, and the fact that stationarity is downward absolute from $V[G]$ to $ult(V, G)$. $\square$

We also state, without proof, a similar characterization for $wDRP(Unif_{\omega_1})$ (here $Unif_{\omega_1}$ denotes the class of ω_1 -sized M such that $cf(\sup(M \cap \lambda)) = \omega_1$ for every $\lambda \in M$ of uncountable cofinality). The proof is very similar to the proof of Theorem 8, except that one would instead use the fact that for $M \in Unif_{\omega_1}$, if $S \in M$ is a stationary subset of $\theta \cap \text{cof}(\omega)$ then S reflects to $\sup(M \cap \theta)$ if and only if V believes that $\sigma_M^{-1}(S)$ is stationary in $\sigma_M^{-1}(\theta)$.

Theorem 10. *For regular $\theta \geq \omega_2$, let Z^θ denote the collection of $M \in \wp_{\omega_2}(H_{(\theta^\omega)^+})$ such that $M \cap H_\theta \in Unif_{\omega_1}$. The following are equivalent:*

- (A) $wDRP(\theta, Z^\theta)$
- (B) *There are stationarily many $M \in Z^\theta$ such that $NS \upharpoonright \theta \cap \text{cof}(\omega)$ condenses correctly via M*
- (C) *There is a normal ideal I whose dual contains the club filter over Z^θ such that $\Vdash_{\mathbb{P}_I}$ “every stationary $R \subset \theta \cap \text{cof}(\omega)$ from the ground model remains stationary in $ult(V, \dot{G})$ ”;*
- (D) *There is a stationary $S \subset Z^\theta$ such that $S \Vdash_{NS}$ “every stationary $R \subset \theta \cap \text{cof}(\omega)$ from the ground model remains stationary in $ult(V, \dot{G})$.”*

4. MODELS OF $DRP(\theta)$ WITH IDEAL FORCINGS THAT HAVE A σ -CLOSED DENSE SUBFORCING

In [10] it was noted that if κ is supercompact, then $MA^{+\omega_1}(\sigma\text{-closed})$ holds in $V^{Col(\omega_1, < \kappa)}$.¹⁰ In that model, a special case of Foreman's Duality Theorem ([6]) implies that for every σ -closed poset $\mathbb{Q}$ and every λ , there is an ideal which concentrates on $S_{\mathbb{Q}} \cap \wp_{\omega_2}(H_\lambda)$ and such that the forcing with the ideal is equivalent to a σ -closed forcing. We give a rough sketch of the argument; more details and similar arguments appear in Section 5, where we obtain analogous results for PFA (but starting from much larger cardinals than a supercompact).

Let κ be supercompact, $\mathbb{P} := Col(\omega_1, < \kappa)$, and G be $(V, \mathbb{P})$ -generic. Let $\mathbb{Q}$ be a σ -closed forcing in $V[G]$, and $\dot{\mathbb{Q}}$ a name for it. Inside $V[G]$, pick a λ sufficiently large such that $Col(\omega_1, \lambda)$ is forcing equivalent to $\mathbb{Q} \times Col(\omega_1, \lambda)$ (see Section 14 of [5]). Note V and $V[G]$ compute $Col(\omega_1, \lambda)$ the same way since they have the same ω sequences. Let $j : V \rightarrow M$ be a H_λ -supercompact ultrapower of V (i.e. the ultrapower map by some normal fine measure on $\wp_\kappa(H_\lambda)$). Since $\mathbb{P}$ has the κ -cc in V then $j \restriction \mathbb{P} : \mathbb{P} \rightarrow j(\mathbb{P})$ is a regular embedding, so the construction in Section 2.4 is applicable; inside $V[G]$ let $F(j)$ be as in Definition 7.

Since $Col(\omega_1, \lambda)$ is forcing equivalent to $\mathbb{Q} \times Col(\omega_1, \lambda)$ and $M[G]$ is closed under λ sequences from $V[G]$, then $M[G]$ sees the forcing equivalence. So $M[G][H]$ always has a $(H_\lambda^{V[G]} = H_\lambda^{M[G]}, \mathbb{Q})$ -generic; call it g . Then g generic clearly transfers via $\hat{j} \restriction H_\lambda^{V[G]}$ to a $(\hat{j}'' H_\lambda^{V[G]}, \hat{j}(\mathbb{Q}))$ -generic $\tilde{g}$, and moreover $\hat{j} \restriction H_\lambda^{V[G]}$ is an element of $M[G][H]$. Thus $M[G][H]$ always believes there is a $(\hat{j}'' H_\lambda^{V[G]}, \hat{j}(\mathbb{Q}))$ -generic, and so by definition of $F(j)$ we have $V[G] \models S_{\mathbb{Q}} \in F(j)$.

Furthermore, suppose $\dot{S} \in \hat{j}'' H_\lambda^{V[G]}$ is a $\hat{j}(\mathbb{Q})$ -name for a stationary subset of ω_1 ; say $\dot{S} = \hat{j}(\dot{\dot{S}})$. Then $H_\lambda[G][g] = M_\lambda[G][g] \models \text{“}\dot{S}_g \text{ is stationary.”}$ Since λ is sufficiently large with respect to $\mathbb{Q}$, $M[G][g] \models \text{“}\dot{S}_g \text{ is stationary”}$ and thus so does $M[G][H]$, since $M[G][H]$ is a stationary set-preserving forcing extension of $M[G][g]$. From this it follows that $MA^{+Diag}(\sigma\text{-closed})$ holds in $V[G]$.

By (7), in $V[G]$ forcing with $F(j)$ is equivalent to forcing with $Col(\omega_1, < j(\kappa))/G$. In particular, the forcing associated with $F(j)$ has a σ -closed dense subset. If the arbitrary $\mathbb{Q}$ we chose at the beginning was, say, $Col(\omega_1, H_\theta)$ for some regular θ then $F(j)$ concentrates on the set $S_{\mathbb{Q}}^{+\omega_1}$, which consists of sets which witness $DRP(\theta)$ structures.

¹⁰Here $Col(\mu, \theta)$ denotes the usual Levy collapse to add a surjection from μ onto θ as in (15.18) on page 238 of Jech [11]. $Col(\mu, < \theta)$ denotes the version which adds surjections from μ to η for every $\eta < \theta$, as in (15.19) on page 238 of Jech [11].

Finally, we observe that this model satisfies “ideal projections as forcing projections” (see Section 2.1) for a pair of ideals whose forcings are σ -closed. Suppose $\kappa < \lambda << \lambda'$ and U is a normal measure on $\wp_\kappa(\lambda')$ and $j_U : V \rightarrow M_U$ the ultrapower map. Let $proj(U)$ be the projection of U to $\wp_\kappa(\lambda)$ and $j_{proj(U)} : V \rightarrow M_{proj(U)}$ the ultrapower map. In $V[G]$ consider the filters $F(U)$ and $F(proj(U))$ (see Definition 7). We need to show $FP(F(U), F(proj(U)))$ holds in $V[G]$. The simple but key observation is that $Col^{M_{proj(U)}}(\omega_1, < j_{proj(U)}(\kappa))$ is a complete subforcing *in the sense of* $V[G]$ of $Col^{M_U}(\omega_1, < j_U(\kappa))$, since all the listed models are closed under ω -sequences (so they all compute the same relevant Levy Collapse posets), and $Col(\omega_1, < j_{proj(U)}(\kappa))$ is a regular subforcing of $Col(\omega_1, < j_U(\kappa))$. This ensures that the canonical map $(F')^+ \rightarrow F^+$ given by $A \mapsto \{Z \cap \lambda \mid Z \in A\}$ is a forcing projection. We omit the details here, but a similar argument is given in Section 6. The point is that the ideal projection from $(F(U))^+ \rightarrow (F(proj(U)))^+$ is the same as the composition of the following sequence of *forcing* projections (it is important that the map numbered 2 in this list is a forcing projection in the sense of $V[G]$); here $\mathbb{P} := Col(\omega_1, < \kappa)$ and $\mathbb{P}'$, $\mathbb{P}''$ are $j_U(\mathbb{P})$ and $j_{proj(U)}(\mathbb{P})$, respectively:

- (1) The dense embedding from $(F(U))^+ \rightarrow ro(\mathbb{P}''/G)$ from (7), given by $A \mapsto \|\check{A} \in \tilde{U}^{\check{H}''}\|_{ro(\mathbb{P}''/G)}$;
- (2) The forcing projection from $ro(\mathbb{P}''/G) \rightarrow ro(\mathbb{P}'/G)$ obtained from the forcing projection $\mathbb{P}''/G \rightarrow \mathbb{P}'/G$ (i.e. obtained via the map which restricts conditions in $\mathbb{P}''$ to their support on $j_{proj(U)}(\kappa)$; recall this is indeed a forcing projection in the sense of $V[G]$ by the remarks above);
- (3) The isomorphism from (a dense subset of) $ro(\mathbb{P}'/G) \rightarrow (F(proj(U)))^+$ obtained from (7) (i.e. the inverse of the map $B \mapsto \|\check{B} \in proj\tilde{j}(U)^{\check{H}'}\|_{ro(\mathbb{P}'/G)}$);

5. PFA AND PROPER IDEALS ON ω_2

In this section we construct a model of *PFA* which has many ideals I such that $\mathbb{P}_I$ is proper. Since there are known models of *PFA* where stationary reflection for subsets of $\omega_2 \cap \text{cof}(\omega)$ fails (see [1]), and since the existence of an ideal I with completeness ω_2 whose forcing is proper implies such stationary reflection,¹¹ we know that *PFA* does not imply

¹¹This is well-known. If an ideal forcing (where the ideal has, say, completeness ω_2) is proper and G is generic for the forcing, then for any $S \subset \omega_2 \cap \text{cof}(\omega)$ in V , S remains stationary in $V[G]$ and thus $S = j(S) \cap \omega_2$ is stationary in $ult(V, G)$, where $j : V \rightarrow_G ult(V, G)$. So by elementarity, V believes every stationary subset of $\omega_2 \cap \text{cof}(\omega)$ reflects.

the existence of ideals on ω_2 whose forcing is proper. In Section 3, however, we showed that DRP (which follows from $PFA^{+\omega_1}$) implies there are ideals whose associated forcings satisfy a very weak form of properness. In this section we show PFA^{+Diag} is consistent with the existence of many¹² ideals whose associated forcings are proper.

Our model of PFA also exhibits weaker versions of properties which are known to be inconsistent with PFA. Foreman and Magidor [9] proved that, under PFA, there is no presaturated ideal on ω_2 and that $(\omega_3, \omega_2) \rightarrow (\omega_2, \omega_1)$ fails (and that these facts are preserved under mild forcing extensions of PFA). In particular, there is no ideal on ω_2 that has some wellfounded generic ultrapower $j_G : V \rightarrow_G \text{ult}(V, G)$ where $\text{ult}(V, G)$ is closed under sequences of length $< j_G(\omega_2)$ from $V[G]$. The following theorem shows that PFA is consistent with the existence of generic ultrapowers that are closed under sequences of length $j_G(\omega_2)$ from the ground model.

The theorem uses a *superhuge* cardinal:

Definition 11. *Let $n \in \omega$ and κ be a cardinal.*

- κ is n -huge iff there is an elementary $j : V \rightarrow M$ where M is transitive, $\text{crit}(j) = \kappa$, and M is closed under $j^n(\kappa)$ sequences. κ is huge iff it is 1-huge.
- κ is super- n -huge iff for every $\gamma > \kappa$ there is an n -huge embedding with critical point κ such that $j(\kappa) > \gamma$. κ is superhuge iff κ is super-1-huge.

See Kanamori [12] for more information about notions related to hugeness. In particular, κ is n -huge iff there is a κ -complete normal ultrafilter U over some $\wp(\lambda)$ and cardinals $\kappa = \lambda_0 < \lambda_1 < \dots < \lambda_n = \lambda$ such that $\{x \subset \lambda \mid \text{otp}(x \cap \lambda_{i+1}) = \lambda_i\} \in U$ for each $i < n$.

Theorem 12. *Suppose there is a super-2-huge cardinal. Then there is a proper forcing extension which satisfies PFA^{+Diag} and has the following property:*

Let $\mathbb{Q}$ be any proper poset. Then for a proper class of inaccessible λ there is an ideal $I_{\lambda, \mathbb{Q}}$ such that:

- $\check{I}_{\lambda, \mathbb{Q}}$ concentrates on $S_{\mathbb{Q}}^{Diag} \cap \{M \subset H_\lambda \mid \text{otp}(M \cap \lambda) = \omega_2 \text{ and } M \cap \omega_2 \in \omega_2\}$ (i.e. $\check{I}_{\lambda, \mathbb{Q}}$ concentrates on sets in $S_{\mathbb{Q}}^{Diag}$ which witness $(\lambda, \omega_2) \rightarrow (\omega_2, \omega_1)$);
- the forcing associated with $I_{\lambda, \mathbb{Q}}$ is proper;

¹²In the sense that for every proper $\mathbb{Q}$ there are proper class many ideal forcings whose ideals concentrate on $S_{\mathbb{Q}}$.

- whenever $j_G : V \rightarrow_G \text{ult}(V, G)$ is a generic ultrapower by $I_{\lambda, \mathbb{Q}}$, then $\text{ult}(V, G)$ is closed under $\lambda = j_G(\omega_2)$ -sequences from V (though not necessarily from $V[G]$)

Proof. Let κ be a super-2-huge cardinal. We will do the standard construction, but for that we need a Laver function for huge embeddings.¹³

Fact 13. *Let $n \in \omega$ and $n \geq 1$. If κ is super- $(n+1)$ -huge, then there is a Laver function $\text{Lav} : \kappa \rightarrow V_\kappa$ for n -huge embeddings; i.e. for every x and every λ there is an n -huge embedding j such that $\text{cr}(j) = \kappa$, $j(\kappa) \geq \lambda$, and $j(\text{Lav})(\kappa) = x$.*

This is well-known, but we include a sketch of the proof, since there is a technical issue here that does not arise when constructing a Laver function for supercompact embeddings.¹⁴ I also thank the anonymous referee for pointing out an error in the original version.

Proof. Mimic the proof of Theorem 20.21 in [11]. Let us say i is an $(\alpha, \geq \lambda)$ - n -huge embedding iff i is n -huge, $\text{cr}(i) = \alpha$, and $i(\alpha) \geq \lambda$; similarly say i is (α, λ) - n -huge if i is n -huge with critical point α and $i(\alpha) = \lambda$. For any α and any partial function $g : \alpha \rightarrow V_\alpha$, set:

- (9) $\lambda_{\alpha, g} :=$ the least ordinal λ such that for some $x \in V_\lambda$, there is no $(\alpha, \geq \lambda)$ - n -huge embedding i such that $i(g)(\alpha) = x$ (if such a λ exists).

Recursively define a partial function $f^* : \kappa \rightarrow V_\kappa$ by:

Assuming $f^* \upharpoonright \alpha$ has been defined:

- (10)
 - (1) If $\lambda_{\alpha, f^* \upharpoonright \alpha}$ is not defined, then $f^*(\alpha)$ is not defined either;
 - (2) If $\lambda_{\alpha, f^* \upharpoonright \alpha}$ is defined but none of the x witnessing this fact are in V_κ , then again $f^*(\alpha)$ is not defined;
 - (3) Otherwise $f^*(\alpha)$ is defined to be some $x \in V_\kappa$ which witnesses that $\lambda_{\alpha, f^* \upharpoonright \alpha}$ is defined

¹³Corazza [2] considered Laver functions for super-almost-huge embeddings, starting from a super 1-huge cardinal. An argument similar to the one below using only a Laver function for almost huge cardinals would still give proper ideals, but we want our ideals to concentrate on models which witness $(\theta, \omega_2) \twoheadrightarrow (\omega_2, \omega_1)$; this is why we instead use a Laver function for huge embeddings.

¹⁴This technical issue occurs in the proof of Claim 14. The issue is that if U is a (κ, λ) -supercompact ultrafilter and $\mu < \lambda$, then the projection of U to $\wp_\kappa(\mu)$ is a (κ, μ) -supercompact ultrafilter. This need not be the case for ultrafilters witnessing hugeness.

Note that if $f^*(\alpha)$ is defined, say $f^*(\alpha) = x_\alpha$, then $x_\alpha \in V_\kappa$ and there is no $(\alpha, \geq \lambda_{\kappa, f^* \upharpoonright \alpha})$ - n -huge embedding i such that $i(f^* \upharpoonright \alpha)(\alpha) = x_\alpha$.

Suppose for a contradiction that κ is super- $(n+1)$ -huge and there is no Laver function from $\kappa \rightarrow V_\kappa$ for n -huge embeddings. So $\lambda_{\kappa, f}$ is defined for every partial $f : \kappa \rightarrow V_\kappa$, and in particular:

$$(11) \quad \lambda_{\kappa, f^*} \text{ is defined (in } V).$$

By super- $n+1$ -hugeness of κ , there is some $n+1$ -huge embedding $j : V \rightarrow M$ such that $cr(j) = \kappa$ and $j(\kappa) > \lambda_{\kappa, f^*}$.

Claim 14. λ_{κ, f^*}^M and $j(f^*)(\kappa)$ are defined, and $\lambda_{\kappa, f^*}^M \leq \lambda_{\kappa, f^*}^V$.

Proof. By (11) and the definition of λ_{κ, f^*} , there is some $x^* \in V_{\lambda_{\kappa, f^*}}$ which witnesses (in V) that λ_{κ, f^*} is defined. Since $\lambda_{\kappa, f^*} < j(\kappa)$ and M is closed under $j(\kappa)$ sequences, then this x^* is an element of $V_{j(\kappa)}^M$.

Set $\lambda^* := \lambda_{\kappa, f^*}^V$. Note that since f^* maps into V_κ , then $j(f^*) \upharpoonright \kappa = f^*$. We will show that $M \models$ “There is no $(\kappa, \geq \lambda^*)$ - n -huge embedding i such that $i(f^*)(\kappa) = x^*$ ”. If we manage to prove this, then since $x^* \in V_{\lambda^*} = V_{\lambda^*}^M$ and $\lambda^* < j(\kappa)$ (by choice of j), then $j(f^*)(\kappa)$ will indeed be defined (since then x^* is an element of $V_{j(\kappa)}^M$ and thus we would be in the last clause, rather than the middle clause, in the definition (10) of $j(f^*)$ inside M). Moreover, this will also show that $\lambda_{\kappa, f^*}^M \leq \lambda^*$ (by minimality in the definition of the $\lambda_{(-, -)}$ function as defined in M). Note x^* need not be equal to $j(f^*)(\kappa)$, but this is not necessary, as we only are trying to show that $j(f^*)(\kappa)$ is defined.

So suppose for a contradiction that $M \models$ “There is some $(\kappa, \geq \lambda^*)$ - n -huge embedding i such that $i(f^*)(\kappa) = x^*$ ”; the quoted statement is Σ_2 in parameters $\kappa, \lambda^*, f^*, x^*$. Since κ is super- $n+1$ -huge then in particular κ is supercompact in V and $j(\kappa)$ is supercompact in M . Then $V_{j(\kappa)}^M \prec_{\Sigma_2} M$ (see Proposition 22.3 of Kanamori [12]). So $V_{j(\kappa)}^M = V_{j(\kappa)} \models$ “There is some $(\kappa, \geq \lambda^*)$ - n -huge embedding i such that $i(f^*)(\kappa) = x^*$ ”. Then there is some $\eta \in [\lambda^*, j(\kappa))$ and a normal measure $U \in V_{j(\kappa)}$ such that $U \subset \wp(\eta)$ codes this n -huge embedding; but since $j(\kappa)$ is inaccessible in V and $\eta < j(\kappa)$ then U also codes an n -huge embedding from the point of view of V .¹⁵ This contradicts the fact that (in V) there is no $(\kappa, \geq \lambda^*)$ - n -huge embedding i such that $i(f^*)(\kappa) = x^*$. $\square$

So by Claim 14 we may set $\bar{x} := j(f^*)(\kappa)$ and $\bar{\lambda} := \lambda_{\kappa, f^*}^M$. We reach a contradiction by showing that $M \models$ “There is some $(\kappa, \geq \bar{\lambda})$ - n -huge embedding i such that $i(f^*)(\kappa) = \bar{x}$ ”. This is the first point where we

¹⁵and $V \models$ “ i_U is $(\kappa, \geq \eta)$ - n -huge and $i_U(f^*)(\kappa) = x^*$ ”.

use the $n + 1$ -hugeness of j (up to now we have only used 1-hugeness of j). Let U be the n -huge ultrafilter derived from j ; that is, set $\kappa_n := j^n(\kappa)$ and $U := \{A \subset \wp(\kappa_n) \mid j''\kappa_n \in j(A)\}$. By standard arguments, if $i_U : V \rightarrow N$ is the ultrapower, then $i_U(f^*)(\kappa) = j(f^*)(\kappa) = \bar{x}$ and $i_U(\kappa) = j(\kappa) > \lambda_{\kappa, f^*} \geq \bar{\lambda}$. Since M is closed under $j^{n+1}(\kappa)$ -sequences from V , then $U \in M$ and $i_U \restriction M$ is the ultrapower map as computed in M ; so in particular $M \models "i_U(f^*)(\kappa) = \bar{x} \text{ and } i_U(f^*)(\kappa) > \bar{\lambda}"$, which is a contradiction. $\square$

We note that the assumptions of Fact 13 can be weakened a bit: in the proof, we only needed the embedding $j : V \rightarrow M$ to have the property that the n -huge ultrafilter derived from j is an element of M . For this it would suffice to assume κ is “super- $n + 1$ -almost-huge” (or in fact much less, though more than super- n -hugeness seems to be required for the argument).

Let $\mathbb{P}$ be the standard Baumgartner forcing to produce a model of PFA, but using the Laver function from Fact 13. So $\mathbb{P}$ is a countable support iteration of length κ where for $\alpha < \kappa$, the α -th component is the $\mathbb{P} \restriction \alpha$ -th evaluation of $Lav(\alpha)$, if $\mathbb{P} \restriction \alpha$ forces $Lav(\alpha)$ to be a proper forcing (see [11] for details). Let G be $(V, \mathbb{P})$ generic. Let $\mathbb{Q} \in V[G]$ be a proper poset, λ a (regular) cardinal, $\dot{Q} \in V$ a name for $\mathbb{Q}$, and $j : V \rightarrow N$ a huge embedding with critical point κ such that $j(\kappa) > \lambda$ and $\dot{Q} = j(Lav)(\kappa)$.

In $V[G]$ let F_j be as in Definition 7. We need to show:

- (1) Forcing with $(F_j)^+$ is a proper forcing;
- (2) Forcing with $(F_j)^+$ yields a generic ultrapower which is closed under $\pi(\omega_2)$ sequences from the ground model, where π is the generic ultrapower embedding;
- (3) F_j concentrates on $S_{\mathbb{Q}}^{Diag}$.

Now N believes $j(\mathbb{P})$ is a countable support iteration of proper forcings, where each forcing in the iteration has size $< j(\kappa)$. Since N is closed under $< j(\kappa)$ sequences, then for every $\xi < j(\kappa)$, N is correct whenever it believes that $j(\mathbb{P}) \restriction \xi$ forces $j(Lav)(\xi)$ to be proper; also N correctly computes the countable support iteration of $j(Lav)$.¹⁶ Thus $V \models "j(\mathbb{P}) \text{ is a countable support iteration of proper forcings and is thus proper.}"$ Then we use the following fact:¹⁷

¹⁶Note these facts only use the *almost* hugeness of j .

¹⁷The author is not aware of a reference for this fact, but it is apparently widely known. One way to prove it is to use the fact that $\mathbb{P}_\alpha$ has the ω -covering property to show that $V^{\mathbb{P}_\alpha}$ believes: “the tail forcing $\mathbb{P}/G_\alpha$ is forcing equivalent to my countable support iteration of the same iterands.”

Fact 15. *If $\mathbb{R}$ is a countable support iteration of proper forcings, then for every $\alpha < lh(\mathbb{R})$: $\mathbb{R}_\alpha \Vdash \text{“}\mathbb{R}/\dot{G}_\alpha \text{ is proper.”}$*

Then with $j(\mathbb{P})$ playing the role of the $\mathbb{R}$ from Fact 15, we have in $V[G]$:

$$(12) \quad j(\mathbb{P})/(j''G) = j(\mathbb{P})/G \text{ is proper}$$

Then (7) on page 8 and (12) on page 17 imply that $\mathbb{P}_{F_j}$ is proper in $V[G]$.

We want to see that $S_{\mathbb{Q}}^{Diag} \in F_j$ (this part only requires the supercompactness of κ and is just the standard argument due to Baumgartner). Let H be any $(V[G], j(\mathbb{P})/G)$ -generic. $V[G]$ and $N[G]$ have sufficient agreement so that $N[G] \models \text{“}\mathbb{Q} \text{ is a proper forcing.”}$ So by the definition of Baumgartner’s forcing, the κ -th component of H is generic for $(N[G], \mathbb{Q})$ and equivalently, generic for $(V[G], \mathbb{Q})$. Let g denote this component. Now $\hat{j}^* := \hat{j}_{G*H} \restriction H_\lambda^{V[G]} \in N[G][H]$; this is standard (it is the inverse of the collapsing map obtained from $j''H_\lambda^V$ and $j''G$). Thus $N[G][H]$ sees that g can be transferred to a $(\hat{j}_{G*H}''H_\lambda[G], \hat{j}_{G*H}(\mathbb{Q}))$ -generic; call this generic $\tilde{g}$. In addition, if $\dot{S} = \hat{j}_{G*H}(\dot{S})$ is any element of $\hat{j}_{G*H}''H_\lambda[G]$ which is a $\hat{j}_{G*H}(\mathbb{Q})$ -name for a stationary subset of ω_1 , then $\dot{S}_g$ is stationary in $H_\lambda^V[G][g] = H_\lambda^N[G][g]$. Since λ is sufficiently large with respect to $\mathbb{Q}$ this implies $\dot{S}_g$ is stationary in $N[G][g]$, and since $N[G][H]$ is a stationary-set-preserving forcing extension of $N[G][g]$ (by Fact 15), $\dot{S}_g$ is still stationary in $N[G][H]$.

Thus we have shown that for arbitrary H , the model $\hat{j}_{G*H}''H_\lambda[G]$ is an element of $\hat{j}_{G*H}(S_{\mathbb{Q}}^{Diag})$; so $S_{\mathbb{Q}}^{Diag} \in F_j$ by the definition of F_j . Also note that since $j''\lambda \in N$ and has ordertype $\lambda = j(\kappa) = \aleph_2^{N[G][H]}$, then $N[G][H] \models otp(\hat{j}_{G*H}''H_\lambda[G] \cap ORD) = \aleph_2$ and $\hat{j}_{G*H}''H_\lambda[G] \cap \aleph_2 \in \aleph_2$. So F_j concentrates on sets which witness $(\lambda, \omega_2) \rightarrow (\omega_2, \omega_1)$.

The closure of the generic ultrapower is proved as follows: say $f : \eta \rightarrow ORD$ is a function in $V[G]$ where $\eta \leq i_K(\omega_2)$ and K is any generic for forcing with F_j^+ over $V[G]$, and i_K is the generic ultrapower map. By (7) there is some H such that $i_K = \hat{j}_{G*H}$ and $\hat{j}_{G*H} : V[G] \rightarrow N[G][H]$ is the generic ultrapower. Say $f = \dot{f}_G$, and for each $\xi < \eta$ let A_ξ be a maximal antichain of conditions in $\mathbb{P}$ which decide the value of $\dot{f}(\check{\xi})$; note each $A_\xi \in V_\kappa \subset N$, so by the hugeness of j we have that $\langle A_\xi | \xi < \eta \rangle$ is an element of N . Thus f is an element of $N[G]$. □

6. PFA PLUS “IDEAL PROJECTIONS AS FORCING PROJECTIONS”

Finally, we prove that if we start with a super-3-huge cardinal, the ideals in Theorem 12 can instantiate very special forms of “Ideal projections as forcing projections”; this is a property which falls somewhere between precipitousness and saturation; moreover, certain instances of “Ideal projections as forcing projections” are *equivalent* to saturation of the projection ideal (see [3]). In particular, we obtain a model of PFA where $FP(I', I)$ for some I' whose dual concentrates on $[\Omega(I)]^\omega$ -faithful models;¹⁸ this property is stronger than properness of $\mathbb{P}_I$.

Suppose that κ is super-3-huge (see Definition 11). By Fact 13, there is a Laver function Lav for 2-huge embeddings; i.e. for every x and every λ there is a $\nu \geq \lambda$ and a 2-huge (κ, ν) -embedding i such that $i(Lav)(\kappa) = x$. Now use this Laver function in the Baumgartner forcing $\mathbb{P}$ for PFA, as in the proof of Theorem 12. Now consider some normal ultrafilter U on $\{X \subset \kappa'' \mid otp(X) = \kappa' \text{ and } otp(X \cap \kappa') = \kappa\}$; note that U gives rise to a 2-huge embedding. Let $j_U : V \rightarrow M_U$; note $j_U(\kappa) = \kappa'$ and M_U is closed under $j_U^2(\kappa) = \kappa''$ sequences.

Let $proj(U)$ be the projection of U to a (κ, κ') -huge ultrafilter,¹⁹ and $j_{proj(U)} : V \rightarrow_{proj(U)} M_{proj(U)}$. Then j_U factors as $k \circ j_{proj(U)}$ such that $k \restriction \kappa' + 1 = id$. Let $\mathbb{P}' := j_{proj(U)}(\mathbb{P})$. Note:

$$(13) \quad j_U(\mathbb{P}) = j_{proj(U)}(\mathbb{P}) = \mathbb{P}'$$

Let G be $(V, \mathbb{P})$ -generic. In $V[G]$ let $\dot{U}''$ denote the $\mathbb{P}'/G$ -name for the $V[G]$ -ultrafilter on $[\kappa'']^{\kappa'}$ derived from $(\hat{j}_U)_H$ (see Section 2.4); and let $\dot{U}'$ denote the $\mathbb{P}'/G$ -name for the $V[G]$ -ultrafilter on $[\kappa']^\kappa$ derived from $(\hat{j}_{proj(U)})_H$ (where H is $(V[G], \mathbb{P}'/G)$ -generic). Let $F'' := F(U)$ and $F' := F(proj(U))$ be as in Definition 7.

Consider the map on $(F'')^+$ defined by $A \mapsto proj(A, \kappa') := \{Z \cap \kappa' \mid Z \in A\}$. We want to see that this is the ideal projection of F'' onto F' , and that this is also a projection in the sense of forcing. So assume $\mathcal{A}$ is a maximal antichain in $(F')^+$ and $A \in (F'')^+$. We need to show there is some $B \in \mathcal{A}$ such that A has F'' -positive intersection with $Lift(B) := \{Z \in [\kappa'']^{\kappa'} \mid Z \cap \kappa' \in B\}$.

Let H be $(V[G], \mathbb{P}'/G)$ -generic such that $A \in \dot{U}''_H$ (such a generic exists since $A \in (F'')^+$, by the definition of F''). Then (note k fixes

¹⁸Here $\Omega(I)$ is some regular cardinal sufficiently large so that the properness, precipitousness, etc. of I^+ is correctly decided by $H_{\Omega(I)}$. Clearly if there is an inaccessible $\mu > trcl(I)$ then $\Omega(I)$ can be taken to be strictly less than the least such μ ; the precise value of $\Omega(I)$ is not important in present application.

¹⁹i.e. $proj(U) = \{\{X \cap \kappa' \mid X \in A\} \mid A \in U\}$.

$\mathbb{P}'$) k can be extended to $\tilde{k} : M_{proj(U)}[G][H] \rightarrow M_U[G][H]$ and $\hat{j}_U^{G*H} = \tilde{k} \circ \hat{j}_{proj(U)}^{G*H}$; and the map $\hat{j}_{proj(U)}^{G*H}$ is (from the point of view of $V[G][H]$) the ultrapower of $V[G]$ by the projection of $\dot{U}''_H$ to $[\kappa']^\kappa$. By (7), $(F')^+$ is forcing equivalent (in $V[G]$) to $\mathbb{P}'$ and so this ultrapower is a generic ultrapower of $V[G]$ by $(F')^+$; so in particular the ultrafilter $proj(\dot{U}''_H)$ meets $\mathcal{A}$; say B is the element of $\mathcal{A}$ which is in $proj(\dot{U}''_H)$. Let $B'' \in \dot{U}''_H$ be such that $B = proj(B'')$. Then $B'' \cap A$ is F'' -positive. This completes the proof that $FP(F'', F')$ holds.

Finally, by the same argument as in the proof of Theorem 12, the almost hugeness of the embeddings and the fact that $\mathbb{P}$ is a countable support iteration of proper forcings ensures that $\mathbb{P}'$ is proper from the point of view of $V[G]$. Moreover we have that $(F'')^+$ and $(F')^+$ are both isomorphic to a dense subset of $\mathbb{P}'$. In particular, forcing with $(F'')^+$ is proper and so F'' must concentrate on $[\Omega(F')]^\omega$ -faithful (in fact $[ORD]^\omega$ -faithful) models.

REFERENCES

- [1] Robert E. Beaudoin, *The proper forcing axiom and stationary set reflection*, Pacific J. Math. **149** (1991), no. 1, 13–24. MR1099782 (92b:03054)
- [2] Paul Corazza, *Laver sequences for extendible and super-almost-huge cardinals*, J. Symbolic Logic **64** (1999), no. 3, 963–983, DOI 10.2307/2586614. MR1779746 (2001g:03096)
- [3] Sean Cox, *Ideal projections as forcing projections*, in preparation.
- [4] ———, *The Diagonal Reflection Principle*, to appear in the Proceedings of the American Mathematical Society.
- [5] James Cummings, *Iterated forcing and elementary embeddings*, Handbook of set theory. Vols. 1, 2, 3, Springer, Dordrecht, 2010, pp. 775–883. MR2768691
- [6] Matthew Foreman, *Calculating quotient algebras for generic embeddings, to appear in Israel J. Math.*
- [7] ———, *Ideals and Generic Elementary Embeddings*, Handbook of Set Theory, Springer, 2010.
- [8] ———, *Smoke and mirrors: combinatorial properties of small cardinals equiconsistent with huge cardinals*, Adv. Math. **222** (2009), no. 2, 565–595, DOI 10.1016/j.aim.2009.05.006. MR2538021
- [9] Matthew Foreman and Menachem Magidor, *Large cardinals and definable counterexamples to the continuum hypothesis*, Ann. Pure Appl. Logic **76** (1995), no. 1, 47–97. MR1359154 (96k:03124)
- [10] M. Foreman, M. Magidor, and S. Shelah, *Martin’s maximum, saturated ideals, and nonregular ultrafilters. I*, Ann. of Math. (2) **127** (1988), no. 1, 1–47. MR924672 (89f:03043)
- [11] Thomas Jech, *Set theory*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2003. The third millennium edition, revised and expanded. MR1940513 (2004g:03071)

- [12] Akihiro Kanamori, *The higher infinite*, 2nd ed., Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2003. Large cardinals in set theory from their beginnings. MR1994835 (2004f:03092)
- [13] Matteo Viale, *Guessing models and generalized Laver diamond (to appear in Annals of Pure and Applied Logic)* (2010).
- [14] W. Hugh Woodin, *The axiom of determinacy, forcing axioms, and the nonstationary ideal*, de Gruyter Series in Logic and its Applications, vol. 1, Walter de Gruyter & Co., Berlin, 1999. MR1713438 (2001e:03001)

E-mail address: `sean.cox@uni-muenster.de`

INSTITUT FÜR MATHEMATISCHE LOGIK UND GRUNDLAGENFORSCHUNG, UNIVERSITÄT MÜNSTER, EINSTEINSTRASSE 62, 48149 MÜNSTER, TEL.: +49-251-83-33 790, FAX: +49-251-83-33 078